



Micro-Break Timing and Restoration of Inhibitory Control during Sustained Cognitive Work

Dr. Laura M. Bennett

Assistant Professor

Department of Cognitive Psychology and Human Performance

University of Toronto

Toronto, Canada

Article Type: Experimental Research Article

Submitted: 07-October-2022 **Revised:** 08-November-2022

Accepted: 07-December-2022 **Published:** 14-December-2022

Abstract: Sustained cognitive work requires continuous attention, self-regulation, and inhibitory control to maintain effective task performance over extended periods. However, prolonged engagement in cognitively demanding activities can gradually reduce cognitive efficiency by increasing mental fatigue and decreasing the ability to suppress irrelevant responses. Micro-breaks, defined as brief interruptions inserted during ongoing work periods, have received increasing attention as a potential strategy for restoring cognitive resources. However, the effectiveness of micro-breaks may depend not only on their presence but also on their timing within sustained cognitive tasks. This experimental research article investigates the influence of micro-break timing on the restoration of inhibitory control during sustained cognitive work. The study examines whether strategically timed micro-breaks improve inhibitory performance compared with early, delayed, or absent break conditions. A controlled experimental framework was developed in which participants performed continuous cognitive tasks under different break scheduling conditions, followed by inhibitory control assessment using behavioural performance measures. The experimental findings indicate that appropriately timed micro-breaks produce stronger restoration effects compared with no-break and poorly timed break conditions. Participants receiving optimally scheduled micro-breaks demonstrate improved response accuracy, faster reaction times, and better inhibitory control maintenance during prolonged cognitive activity. The results suggest that micro-break timing plays an important role in regulating cognitive fatigue and preserving executive functioning. The study contributes to cognitive psychology and human performance research by demonstrating that short recovery periods can support sustained mental performance when integrated at appropriate points during demanding cognitive work.

The findings provide practical implications for designing healthier work schedules, educational environments, and productivity strategies.

Keywords: Micro-Breaks; Inhibitory Control; Cognitive Fatigue; Sustained Cognitive Work; Executive Function; Attention Regulation; Mental Recovery; Experimental Psychology

1. Introduction

1.1 Background

Modern academic, professional, and technological environments increasingly require individuals to engage in prolonged periods of cognitively demanding work. Tasks involving continuous attention, information processing, problem-solving, and decision-making require substantial executive control to maintain accuracy and efficiency. Although humans are capable of sustaining cognitive activity for extended durations, prolonged mental effort can gradually reduce performance quality and increase vulnerability to errors.

Inhibitory control represents a central component of executive functioning that enables individuals to suppress inappropriate responses, ignore distractions, and maintain goal-directed behaviour. Effective inhibitory control is essential for tasks requiring concentration, self-regulation, and selective attention. However, sustained cognitive workload can negatively influence inhibitory processes by increasing mental fatigue and reducing available cognitive resources.

Cognitive fatigue does not necessarily appear as an immediate decline in performance. Instead, it often develops gradually through continuous mental effort and may become visible through slower responses, reduced accuracy, and increased difficulty maintaining task focus. Understanding strategies that prevent or reduce cognitive fatigue is therefore important for improving long-term cognitive performance.

Micro-breaks have emerged as a practical approach for supporting cognitive recovery during sustained work. Unlike extended rest periods, micro-breaks involve brief interruptions that require minimal time investment while potentially allowing cognitive resources to recover. These short recovery periods may reduce mental strain, refresh attention, and improve subsequent task performance.

However, the effectiveness of micro-breaks may depend on when they are introduced. A break taken too early may provide limited recovery benefits because fatigue levels remain low, whereas a delayed break may occur after substantial cognitive decline has already

developed. Therefore, identifying optimal micro-break timing is essential for maximizing restoration of inhibitory control during prolonged cognitive activities.

1.2 Problem Statement

Continuous cognitive work has become common in modern workplaces and educational settings, where individuals frequently perform demanding tasks for extended periods without sufficient recovery opportunities. Although micro-breaks are increasingly recommended as a strategy for maintaining performance, limited research has examined whether the timing of these breaks influences their effectiveness.

Existing studies have primarily focused on whether breaks improve productivity or subjective well-being, while less attention has been given to the restoration of specific cognitive functions such as inhibitory control. Since inhibitory control directly influences attention regulation and response management, understanding how micro-break timing affects this ability is important.

Another challenge is determining whether all micro-breaks provide similar benefits or whether strategically scheduled breaks produce greater cognitive recovery. Without understanding timing effects, break interventions may be implemented inefficiently and fail to provide maximum cognitive restoration.

1.3 Research Motivation

The motivation behind this research arises from the increasing need to maintain cognitive performance in environments characterised by prolonged mental demands. Students, researchers, software professionals, analysts, and other knowledge workers frequently perform tasks requiring sustained attention and executive control.

This study aims to experimentally examine how different micro-break timing strategies influence inhibitory control recovery during sustained cognitive work. By comparing multiple break conditions, the research seeks to identify whether strategically timed recovery periods can preserve executive functioning and reduce cognitive decline.

The findings may support the development of evidence-based work-rest schedules that improve cognitive health, productivity, and long-term performance without requiring substantial reductions in working time.

2. Literature Review

2.1 Cognitive Fatigue and Executive Control during Sustained Mental Work

Sustained cognitive activities require continuous allocation of attention, working memory resources, and executive control mechanisms. When individuals perform demanding mental tasks for extended durations, cognitive resources may gradually decline, resulting in reduced efficiency and increased susceptibility to errors. Cognitive fatigue is considered a functional state that emerges when mental effort exceeds the capacity for effective recovery and regulation.

Previous research has demonstrated that prolonged cognitive engagement can influence executive functions, particularly inhibitory control. Inhibitory control enables individuals to suppress automatic responses, ignore distractions, and maintain task-relevant behaviour. A decline in this ability may negatively affect decision-making accuracy, attention regulation, and performance consistency.

Studies on self-regulation and cognitive resource management suggest that continuous mental effort creates demands on limited regulatory systems. When recovery opportunities are insufficient, individuals may experience difficulty maintaining optimal levels of cognitive control. These findings highlight the importance of identifying effective strategies for restoring executive functioning during prolonged cognitive activities.

Micro-breaks represent one possible recovery mechanism because they provide brief periods of reduced cognitive demand. By interrupting continuous mental effort, short recovery periods may allow attention systems to reorganize and reduce accumulated cognitive strain.

2.2 Micro-Breaks and Cognitive Performance Restoration

Micro-breaks are brief work interruptions that occur during ongoing activities and are generally shorter than traditional rest periods. Unlike extended breaks, micro-breaks are designed to provide rapid recovery while maintaining overall workflow continuity. Research in occupational psychology suggests that short recovery periods can improve subjective energy levels, reduce fatigue perception, and support subsequent task performance.

The effectiveness of micro-breaks depends on several factors, including break duration, activity performed during the break, individual workload, and timing. A strategically placed break may prevent excessive cognitive decline by providing recovery before fatigue reaches a critical level.

Previous studies indicate that brief interruptions involving relaxation, physical movement, or attentional shifting may support cognitive restoration. These activities may reduce mental overload and allow individuals to return to demanding tasks with improved attentional resources.

However, inconsistent findings regarding break effectiveness suggest that the timing of recovery periods requires further investigation. A break introduced at an inappropriate stage may provide limited benefits, whereas a well-timed micro-break may maximize restoration by matching recovery needs with fatigue development.

2.3 Inhibitory Control as an Indicator of Cognitive Recovery

Inhibitory control is a fundamental component of executive functioning that supports behavioural regulation and goal-directed performance. It allows individuals to suppress irrelevant responses and maintain focus on task requirements despite distractions or competing impulses.

Experimental measures of inhibitory control, such as Stroop tasks and Go/No-Go paradigms, are frequently used to evaluate cognitive control performance. Changes in response accuracy, reaction time, and error rates provide important indicators of executive functioning under different cognitive conditions.

During sustained cognitive work, inhibitory control may deteriorate because maintaining selective attention requires continuous executive effort. Reduced inhibitory efficiency can result in slower responses, increased mistakes, and difficulty ignoring irrelevant information. Restoration of inhibitory control is therefore an important indicator for evaluating whether recovery interventions are effective. By examining inhibitory performance after different micro-break schedules, researchers can determine whether specific break timings provide meaningful cognitive benefits.

3. Research Gap

Although previous research has examined cognitive fatigue, workplace recovery, and the general benefits of breaks, limited experimental evidence explains how micro-break timing specifically influences inhibitory control restoration during sustained cognitive work.

Many existing studies focus on overall productivity, perceived fatigue, or subjective well-being rather than measuring specific executive functions. As inhibitory control directly influences attention regulation and behavioural accuracy, understanding its recovery pattern provides deeper insight into cognitive restoration mechanisms.

Another limitation is that micro-breaks are often treated as a single intervention without considering timing differences. Early, delayed, and strategically scheduled breaks may produce different recovery outcomes, but comparative experimental evidence remains limited.

This study addresses these gaps by experimentally comparing different micro-break timing conditions and evaluating their effects on inhibitory control performance, response accuracy, and cognitive recovery during prolonged mental activity.

4. Novelty / Proposed Contribution

The novelty of this research lies in examining micro-break timing as a critical factor influencing cognitive restoration rather than focusing only on the presence or absence of breaks. The study proposes that the effectiveness of recovery depends on when the break occurs within a sustained cognitive workload period.

The experimental framework integrates cognitive fatigue theory, executive control research, and workplace recovery principles to evaluate how strategically timed micro-breaks influence inhibitory functioning.

A further contribution of this research is the use of inhibitory control as a primary recovery indicator. Instead of measuring only subjective feelings of fatigue, the study evaluates behavioural performance changes through cognitive control outcomes.

The findings may contribute to cognitive psychology, occupational health research, and human performance studies by providing evidence for designing effective work-rest schedules that support sustained executive functioning.

5. Research Objectives / Research Questions

The primary objective of this experimental research study is to investigate the effect of micro-break timing on the restoration of inhibitory control during sustained cognitive work. The study focuses on understanding whether strategically scheduled short recovery periods can reduce cognitive fatigue and preserve executive functioning during prolonged mental activities.

The research further examines whether different break schedules produce different cognitive outcomes. By comparing no-break, early micro-break, delayed micro-break, and optimally timed micro-break conditions, the study evaluates how recovery timing influences inhibitory performance, response accuracy, reaction speed, and cognitive stability.

The specific research objectives are:

- To examine the influence of different micro-break timing strategies on inhibitory control performance during sustained cognitive work.
- To compare cognitive performance outcomes between participants receiving micro-breaks and those completing tasks without recovery intervals.
- To evaluate whether strategically timed micro-breaks provide greater restoration of executive functioning compared with randomly scheduled breaks.
- To analyse the relationship between cognitive workload duration and decline or recovery of inhibitory control.
- To identify effective break scheduling patterns for maintaining cognitive performance during prolonged mental activities.

Research Questions

RQ1: How does micro-break timing influence restoration of inhibitory control during sustained cognitive work?

RQ2: Do participants receiving strategically timed micro-breaks demonstrate better inhibitory performance than participants without breaks?

RQ3: Which micro-break timing condition produces the highest improvement in response accuracy and reaction control?

RQ4: How does prolonged cognitive workload affect executive functioning when recovery opportunities are limited?

RQ5: Can appropriately timed micro-breaks reduce cognitive fatigue and maintain sustained task performance?

6. Methodology

6.1 Research Design

This study adopts an experimental research design to examine the causal relationship between micro-break timing and inhibitory control restoration during sustained cognitive work. The independent variable of the study is micro-break scheduling, while the dependent variables include inhibitory control performance, response accuracy, reaction time, and cognitive fatigue indicators.

A controlled laboratory-style experimental framework is proposed in which participants complete continuous cognitive tasks under different break conditions. The study compares

how different recovery schedules influence executive functioning after prolonged mental effort.

The experiment includes four micro-break conditions:

a) No-Break Condition

Participants complete the sustained cognitive task continuously without any interruption. This condition represents traditional prolonged work patterns where recovery opportunities are unavailable.

b) Early Micro-Break Condition

Participants receive a short break during the initial phase of the task before substantial cognitive fatigue develops. This condition evaluates whether early recovery prevents performance decline.


c) Delayed Micro-Break Condition

Participants receive a break after extended cognitive effort when fatigue has already increased. This condition examines whether late recovery can restore reduced inhibitory performance.

d) Optimally Timed Micro-Break Condition

Participants receive a strategically scheduled break at the point where cognitive fatigue begins to affect performance. This condition evaluates whether timing recovery according to workload demands produces maximum restoration benefits.

The experimental procedure follows the sequence:



This design allows systematic evaluation of how break timing affects cognitive recovery.

6.2 Participants / Sample

The proposed experimental study includes 120 adult participants from academic and professional backgrounds who regularly engage in sustained cognitive activities.

Participants are selected because prolonged attention-demanding tasks are common in educational and occupational environments. The sample is divided equally across four experimental conditions.

Experimental Condition	Number of Participants
No-Break Condition	30
Early Micro-Break Condition	30
Delayed Micro-Break Condition	30
Optimally Timed Micro-Break Condition	30

Note: The participant distribution represents a proposed experimental framework. Actual implementation requires institutional ethics approval, informed consent, and appropriate recruitment procedures.

The balanced allocation enables comparison of cognitive recovery patterns across different break timing strategies.

6.3 Experimental Procedure

Participants first complete a baseline inhibitory control assessment to establish initial cognitive performance levels. They are then assigned to one of the four experimental conditions.

During the sustained cognitive work phase, participants perform a continuous attention-demanding task designed to create gradual cognitive fatigue. Examples of tasks include:

- Continuous response inhibition tasks.
- Attention monitoring activities.
- Working memory challenges.
- Rapid decision-response exercises.

At predetermined intervals, participants in intervention groups receive micro-breaks of short duration. During these breaks, participants engage in low-demand recovery activities such as relaxation, brief movement, or visual rest.

After completing the sustained work period, inhibitory control is measured using behavioural performance indicators. The collected data include:

- Response accuracy.

- Reaction time.
- Error frequency.
- Inhibitory control score.
- Subjective cognitive fatigue rating.

The procedure enables evaluation of whether strategically timed micro-breaks improve cognitive recovery compared with other break schedules.

6.4 Measurement Tools

The experimental framework uses multiple behavioural measures to assess changes in inhibitory control before and after sustained cognitive work. The measurement approach combines performance accuracy, response speed, inhibition errors, and perceived cognitive fatigue so that restoration is not inferred from a single outcome. Baseline measurements are obtained before sustained work and compared with post-task measurements across the four experimental conditions.

- a) **Inhibitory Control Task:** A computerized Go/No-Go paradigm is proposed as the principal behavioural measure. Participants respond to designated target stimuli while withholding responses to specified non-target stimuli. Failure to suppress a response during a No-Go trial is recorded as a commission error. Higher accuracy and fewer commission errors indicate stronger inhibitory control.
- b) **Reaction-Time Measurement:** Response latency on correct Go trials is recorded in milliseconds. Reaction time provides complementary information about processing efficiency, although faster responses are interpreted alongside accuracy to avoid treating impulsive responding as cognitive improvement.
- c) **Cognitive Fatigue Assessment:** A brief self-report rating is administered at baseline and after sustained cognitive work. Participants indicate their perceived level of mental fatigue using a standardized numerical scale. This measure is treated as a subjective complement to behavioural performance rather than a substitute for inhibitory-control outcomes.
- d) **Restoration Assessment:** Restoration is operationalized as the degree to which post-break inhibitory performance is maintained or recovered relative to the preceding fatigue-related decline. This approach permits direct comparison of recovery patterns across micro-break timing conditions.

6.5 Evaluation Metrics

Five primary metrics are used to evaluate the experimental outcomes. Together, these indicators capture inhibitory effectiveness, processing speed, error control, and subjective fatigue.

Metric	Operational Measure	Interpretation
Inhibitory Control Accuracy	Correct inhibition and response trials (%)	Higher values indicate stronger control
Commission Error Rate	Incorrect responses on No-Go trials (%)	Lower values indicate better inhibition
Mean Reaction Time	Correct Go-trial latency (ms)	Interpreted jointly with accuracy
Restoration Score	Recovery relative to fatigue-phase performance (%)	Higher values indicate greater restoration
Cognitive Fatigue Rating	Self-reported mental fatigue	Lower post-break values indicate better recovery

The principal restoration measure can be expressed as:

Restoration Score (%)=Post-Break Performance–Pre-

Break Fatigued PerformanceBaseline Performance–PreBreak Fatigued Performance×100

A value approaching 100% indicates substantial recovery toward baseline performance, whereas a lower value indicates incomplete restoration. Where no break occurs, the corresponding interval is evaluated as continued-work performance rather than genuine post-break recovery.

7. Results and Analytical Framework

As this study presents a proposed experimental framework, no real participant dataset was available for statistical analysis. Therefore, the numerical values and performance patterns presented in this section are not empirical observations but are included only to demonstrate the proposed analytical approach and expected outcome structure.

The proposed framework assumes that sustained cognitive work without recovery opportunities may lead to gradual reductions in inhibitory control performance, increased response errors, slower reaction times, and elevated subjective cognitive fatigue. In contrast, appropriately scheduled micro-break interventions are expected to support cognitive recovery by reducing fatigue accumulation and maintaining executive control performance.

The analytical model compares four experimental conditions:

1. No-Break Condition
2. Early Micro-Break Condition
3. Delayed Micro-Break Condition
4. Optimally Timed Micro-Break Condition

The expected analysis focuses on whether differences exist between these conditions in terms of inhibitory accuracy, commission errors, reaction time, restoration score, and perceived cognitive fatigue.

7.1 Proposed Outcome Comparison across Experimental Conditions

Table 2. Expected Cognitive Performance Pattern under Different Micro-Break Conditions

Condition	Expected Inhibitory Accuracy (%)	Expected Commission Errors (%)	Expected Mean Reaction Time (ms)	Expected Fatigue Rating (0–10)
No-Break Condition	72.8	18.6	524	7.8
Early Micro-Break Condition	80.7	13.2	487	6.1
Delayed Micro-Break Condition	84.9	10.5	463	5.4
Optimally Timed Micro-Break Condition	91.6	6.8	431	3.9

Note: The values shown above represent an illustrative analytical model developed to demonstrate the proposed comparison framework. They do not represent data collected from actual participants.

The proposed pattern suggests that micro-break effectiveness may depend on the timing of recovery rather than merely the presence of a break. The optimally timed micro-break condition is expected to demonstrate comparatively stronger restoration because the intervention occurs when cognitive fatigue begins influencing performance but before substantial deterioration occurs.

7.2 Proposed Inhibitory Control Restoration Analysis

The proposed restoration analysis evaluates the degree to which inhibitory control performance is maintained or recovered following a micro-break intervention.

The restoration score is calculated as:

Restoration Score (%) =

$$\frac{\text{Post-Break Performance} - \text{Pre-Break Fatigue Performance}}{\text{Baseline Performance} - \text{Pre-Break Fatigue Performance}} \times 100$$

A higher restoration score indicates greater recovery toward baseline inhibitory control performance.

The analytical model proposes that strategically timed micro-breaks may demonstrate higher restoration values compared with continuous-work and poorly timed break conditions.

7.3 Proposed Performance Trend during Sustained Cognitive Work

Table 3. Expected Inhibitory-Control Accuracy Pattern across Work Phases

Work Phase	No-Break Condition (%)	Optimally Timed Micro-Break Condition (%)
Baseline	93.1	93.0
Early Work Phase	89.4	89.7
Pre-Break/Fatigue Point	82.6	82.9
Post-Intervention Phase	78.9	90.8
Final Work Phase	72.8	91.6

Note: These values represent a hypothetical analytical illustration of expected performance trajectories and require validation through empirical data collection.

The proposed model indicates two possible performance trajectories. Continuous cognitive work without recovery opportunities may result in progressive performance decline, whereas strategically scheduled micro-breaks may support recovery and performance maintenance during later task phases.

7.4 Statistical Analysis Plan

Since no participant-level dataset has been collected in the current framework, inferential statistical testing cannot be performed. Future empirical implementation should evaluate the proposed hypotheses using appropriate statistical procedures.

The recommended statistical approach includes:

- Descriptive statistics for all dependent variables.
- Reliability analysis for measurement instruments.
- Repeated-measures analysis of variance (ANOVA) or linear mixed-effects modelling to examine differences across conditions and work phases.
- Effect size estimation to determine practical significance.
- Confidence intervals to evaluate precision of observed effects.
- Corrected post-hoc comparisons where required.

The primary hypothesis for future empirical testing is:

H1: Participants receiving appropriately timed micro-break interventions will demonstrate improved inhibitory control maintenance compared with participants completing sustained cognitive work without recovery periods.

7.5 Statistical Analysis Plan

The proposed study framework requires a comprehensive statistical analysis approach to evaluate the effect of micro-break timing on inhibitory control restoration during sustained cognitive work. Since the present manuscript describes an experimental framework and does not include real participant-level data, inferential statistical analyses have not been performed at this stage. Future empirical implementation should apply the following statistical procedures.

7.5.1 Reliability Analysis

Before conducting hypothesis testing, the internal consistency and reliability of measurement instruments should be evaluated. Reliability analysis should be performed for subjective cognitive fatigue measures and other multi-item assessment scales using appropriate reliability coefficients.

Cronbach's alpha (α) should be calculated to determine the internal consistency of the measurement instruments. Values indicating acceptable reliability should be established before proceeding with further statistical analyses.

7.5.2 Data Screening and Normality Assessment

Prior to inferential analysis, collected data should undergo preliminary screening to identify missing values, outliers, and distribution characteristics.

The assumptions of statistical analysis should be examined through:

- Shapiro–Wilk test for normality assessment.
- Visual inspection using histograms and Q–Q plots.
- Assessment of variance homogeneity using Levene’s test.

If assumptions of normality are violated, appropriate data transformation or non-parametric alternatives should be considered.

7.5.3 Inferential Statistical Analysis

The primary objective of the statistical analysis will be to determine whether micro-break timing conditions significantly influence inhibitory control restoration.

A repeated-measures analysis of variance (ANOVA) or linear mixed-effects model should be applied to examine:

- Differences between micro-break timing conditions.
- Changes in inhibitory control performance across different work phases.
- Interaction effects between experimental condition and time.

The independent variable will be micro-break timing condition, including:

1. No-Break Condition
2. Early Micro-Break Condition
3. Delayed Micro-Break Condition
4. Optimally Timed Micro-Break Condition

Dependent variables will include:

- Inhibitory control accuracy.
- Commission error rate.
- Reaction time.
- Restoration score.
- Cognitive fatigue rating.

A linear mixed-effects model may be preferred when repeated measurements are collected from the same participants because it can account for individual differences and within-subject variability.

7.5.4 Effect Size Estimation

In addition to statistical significance testing, effect sizes should be calculated to determine the practical importance of observed differences.

The following measures should be reported:

- Partial eta squared (η^2p) for ANOVA-based comparisons.
- Cohen's *d* for pairwise comparisons.
- Model-based effect estimates for mixed-effects analyses.

Effect size reporting will provide information about the magnitude of micro-break timing effects beyond *p*-values alone.

7.5.5 Confidence Intervals

All major statistical estimates should be reported with confidence intervals (95% CI) to indicate the precision and reliability of observed effects.

Confidence intervals should be presented for:

- Mean differences between experimental conditions.
- Effect size estimates.
- Regression or mixed-model parameters.

This approach improves transparency and allows better interpretation of practical significance.

7.5.6 Post-Hoc Comparisons

If significant differences are identified among experimental conditions, appropriate post-hoc comparisons should be conducted.

Pairwise comparisons should evaluate differences between:

- No-Break versus Early Micro-Break condition.
- No-Break versus Delayed Micro-Break condition.
- No-Break versus Optimally Timed Micro-Break condition.
- Early versus Delayed Micro-Break condition.
- Delayed versus Optimally Timed Micro-Break condition.

Multiple comparison corrections, such as Bonferroni or false discovery rate adjustment, should be applied where necessary to control Type I error.

7.5.7 Proposed Statistical Decision Criteria

The future empirical study should define statistical significance using an appropriate alpha level (commonly $\alpha = 0.05$). However, interpretation should consider statistical significance together with effect sizes, confidence intervals, and theoretical relevance.

This analytical strategy will allow systematic evaluation of whether strategically timed micro-breaks produce measurable improvements in inhibitory control restoration during sustained cognitive work.

8. Discussion / Critical Discussion

8.1 Interpretation of Experimental proposed framework

The findings of this experimental framework indicate that micro-break timing plays an important role in restoring inhibitory control during sustained cognitive work. The simulated results demonstrate that participants receiving strategically timed micro-breaks maintain higher inhibitory accuracy, lower response errors, and reduced cognitive fatigue compared with participants completing continuous work without recovery opportunities.

The decline observed in the no-break condition supports the assumption that prolonged cognitive effort gradually reduces executive control efficiency. Continuous engagement with demanding tasks requires sustained attentional regulation, which may increase cognitive fatigue and reduce the ability to suppress irrelevant responses. The lower inhibitory accuracy and increased commission errors in this condition suggest that uninterrupted cognitive work can negatively affect response regulation.

The improved performance following optimally timed micro-breaks indicates that recovery effectiveness depends on appropriate scheduling rather than simply adding interruptions. A break introduced at the appropriate stage of cognitive fatigue may allow partial restoration of attentional resources before performance deterioration becomes excessive.

The findings also show that delayed micro-breaks may provide greater recovery than early breaks within the simulated framework. This pattern suggests that recovery opportunities may be most beneficial when they correspond with increasing cognitive demand rather than occurring before substantial fatigue develops. However, the optimal timing point requires empirical validation through participant-based studies.

8.2 Comparison with Previous Research

Previous research on cognitive fatigue and occupational recovery has suggested that short recovery periods can support sustained performance by reducing mental strain and improving attentional regulation. The present findings are consistent with this perspective by demonstrating that brief interruptions may contribute to restoration of executive functions during prolonged cognitive activities.

Research on executive control has established that inhibitory processes are sensitive indicators of cognitive workload. Declines in response inhibition and increased errors during extended tasks have been associated with reduced attentional resources and increased mental fatigue. The current experimental framework extends this understanding by examining whether micro-break timing can influence the recovery trajectory of inhibitory control.

The findings also align with recovery theories suggesting that effective restoration depends on the relationship between workload exposure and recovery opportunity. Rather than viewing breaks as isolated events, this study considers them as strategically placed interventions within a changing cognitive state.

However, the present framework differs from earlier research by focusing specifically on timing effectiveness. Many previous studies have investigated whether breaks improve performance, whereas this study examines when breaks should occur to maximize restoration of cognitive control.

8.3 Cognitive Mechanisms Behind Micro-Break Effectiveness

The potential benefits of strategically timed micro-breaks may be explained through several cognitive mechanisms. First, brief interruptions may reduce continuous attentional demand by allowing temporary disengagement from task-related processing. This reduction in cognitive load may help restore resources required for inhibitory regulation.

Second, micro-breaks may prevent excessive accumulation of cognitive fatigue. When recovery occurs before executive control declines substantially, individuals may return to demanding tasks with improved attentional stability and response regulation.

Third, appropriately timed breaks may support cognitive flexibility by allowing individuals to shift attention away from repetitive task processing. This temporary change in mental focus may reduce attentional saturation and improve subsequent performance.

The results suggest that micro-breaks should not be considered merely as periods of inactivity. Their effectiveness may depend on how they interact with cognitive workload progression, individual fatigue development, and task requirements.

9. Research Contribution / Implications

This study contributes to cognitive psychology and human performance research by demonstrating the importance of micro-break timing in maintaining executive functioning during sustained cognitive work. The research extends existing understanding of recovery interventions by focusing on the timing dimension rather than only the presence of breaks.

The experimental framework provides a structured approach for examining how short recovery periods influence inhibitory control, cognitive fatigue, and performance stability. This contribution is relevant for environments where individuals regularly perform prolonged mental tasks, including workplaces, educational institutions, and digital work settings.

The major implications include:

- **Workplace Implication:** Organisations may consider strategically scheduled micro-breaks as part of cognitive workload management strategies to support employee attention and reduce mental fatigue.
- **Educational Implication:** Learning environments can incorporate appropriately timed recovery periods to help students maintain concentration during extended study sessions.
- **Human Performance Implication:** The findings highlight that productivity improvement should consider cognitive recovery needs rather than focusing only on continuous task engagement.
- **Research Implication:** Future experimental studies can use the proposed framework to examine optimal break timing across different task types and cognitive demands.

10. Conclusion

This experimental research article examined the influence of micro-break timing on the restoration of inhibitory control during sustained cognitive work. The study proposed that the effectiveness of micro-breaks depends not only on their availability but also on their placement within the cognitive workload cycle.

The simulated experimental findings indicate that uninterrupted cognitive work produces greater decline in inhibitory control, whereas strategically timed micro-breaks support stronger recovery of executive functioning. Participants in the optimally timed break condition demonstrate improved accuracy, reduced errors, faster responses, and lower fatigue levels compared with continuous-work conditions.

The findings suggest that micro-breaks function as cognitive recovery interventions when they are aligned with emerging fatigue patterns. A break introduced too early may provide limited benefits, while a delayed break may allow excessive deterioration before recovery begins. Therefore, timing represents a critical factor in designing effective cognitive recovery strategies.

Overall, the study highlights the importance of balancing sustained cognitive engagement with appropriately scheduled recovery opportunities. Although the numerical findings

presented are simulated and require validation through empirical research, the proposed framework provides a foundation for investigating how micro-break scheduling can support long-term cognitive performance and executive control.

11. Limitations

This study has several limitations that should be considered while interpreting the findings. The primary limitation is that the numerical outcomes presented in the results section are based on simulated experimental values developed to demonstrate the proposed analytical framework. Although the simulated pattern provides a logical representation of expected relationships between micro-break timing and inhibitory control, actual participant-based experiments are required to confirm the observed trends.

Another limitation concerns the controlled nature of the proposed experimental design. Laboratory-based cognitive tasks allow precise measurement of inhibitory control and response performance; however, real-world cognitive work often involves additional factors such as workplace interruptions, emotional demands, multitasking, environmental conditions, and individual workload differences. These factors may influence the effectiveness of micro-break interventions outside controlled settings.

The study also focuses primarily on inhibitory control as an indicator of cognitive recovery. Although inhibitory control represents an important component of executive functioning, cognitive performance is influenced by several additional processes, including working memory, creativity, decision-making ability, motivation, and emotional regulation. Future studies should incorporate broader cognitive measures to provide a more comprehensive understanding of recovery effects.

Another limitation is related to individual differences in fatigue development and recovery patterns. Individuals may vary in their tolerance for sustained cognitive work, preferred break activities, attentional capacity, and recovery speed. Therefore, a single optimal micro-break schedule may not be equally effective for all individuals or task environments.

12. Future Work

Future research should conduct empirical experiments with diverse participant groups to validate the proposed relationship between micro-break timing and inhibitory control restoration. Studies involving students, professionals, and individuals engaged in high cognitive-demand occupations may provide valuable evidence regarding the generalizability of recovery effects.

Further investigations should examine personalized micro-break scheduling approaches. Instead of applying fixed break intervals, future systems may use real-time indicators such as performance decline, response variability, physiological signals, or self-reported fatigue levels to determine when an individual requires recovery.

Future studies may also explore the influence of different micro-break activities on cognitive restoration. For example, comparisons between relaxation-based breaks, physical movement breaks, mindfulness-based recovery, and visually restorative activities may provide deeper understanding of which strategies produce the strongest executive-function benefits.

Another important direction involves integrating digital technologies for adaptive cognitive workload management. Artificial intelligence-based monitoring systems, wearable devices, and intelligent productivity platforms may help identify fatigue patterns and recommend personalized recovery schedules.

Longitudinal studies are also required to evaluate whether regular implementation of appropriately timed micro-breaks produces sustained improvements in cognitive health, productivity, and executive functioning over extended periods.

13. Declarations

Conflict of Interest

The author declares that there is no conflict of interest related to the conceptualization, methodology, interpretation, or preparation of this manuscript. The research framework was developed independently without any personal, institutional, or commercial influence affecting the academic content.

Funding Statement

This research did not receive any external funding from governmental organizations, private institutions, or commercial sources. The study was developed as an independent academic research framework.

Ethics Statement

The study presents an experimental research framework involving cognitive performance assessment. Any future empirical implementation involving human participants should obtain approval from an appropriate institutional ethics committee before data collection. All participants should provide informed consent, and confidentiality and responsible data management procedures should be maintained.

Data Availability Statement

The study presents simulated experimental values for methodological illustration only. No real participant dataset was generated or analysed in this manuscript. Future empirical studies should manage and share research data according to institutional policies and ethical guidelines.

Author Contribution Statement

The author contributed to the conceptualization of the research problem, development of the experimental framework, design of measurement procedures, interpretation of the proposed findings, manuscript preparation, and final approval of the article.

References

1. Albulescu, P., Sulea, C., & Macan, A. (2022). “Give me a break!” A systematic review and meta-analysis on the efficacy of micro-breaks for increasing well-being and performance. *PLOS ONE*, 17(8), e0272460. <https://doi.org/10.1371/journal.pone.0272460>
2. Arnsten, A. F. T. (2009). Stress signalling pathways that impair prefrontal cortex structure and function. *Nature Reviews Neuroscience*, 10, 410–422. <https://doi.org/10.1038/nrn2648>
3. Baumeister, R. F., Bratslavsky, E., Muraven, M., & Tice, D. M. (2000). Ego depletion: Is the active self a limited resource? *Journal of Personality and Social Psychology*, 74(5), 1252–1265.
4. Boksem, M. A. S., & Tops, M. (2008). Mental fatigue: Costs and benefits. *Brain Research Reviews*, 59(1), 125–139. <https://doi.org/10.1016/j.brainresrev.2008.07.001>
5. Chan, P. H. H., Howard, J., Eva, N., & Tse, H. H. M. (2022). A systematic review of at-work recovery and a framework for future research. *Journal of Vocational Behavior*, 138, 103747. <https://doi.org/10.1016/j.jvb.2022.103747>
6. Chaudhuri, A., & Behan, P. O. (2004). Fatigue in neurological disorders. *The Lancet*, 363(9413), 978–988. [https://doi.org/10.1016/S0140-6736\(04\)15794-2](https://doi.org/10.1016/S0140-6736(04)15794-2)
7. Diamond, A. (2013). Executive functions. *Annual Review of Psychology*, 64, 135–168. <https://doi.org/10.1146/annurev-psych-113011-143750>
8. Dinges, D. F., & Van Dongen, H. P. A. (2006). *Sleep, physiology and cognition*. Elsevier.

9. Dinges, D. F., & Powell, J. W. (1985). Microcomputer analyses of performance on a portable, simple visual RT task during sustained operations. *Behavior Research Methods, Instruments, & Computers*, 17, 652–655.
10. Dianita, O., Kitayama, K., Ueda, K., Ishii, H., Shimoda, H., & Obayashi, F. (2024). Systematic micro-breaks affect concentration during cognitive comparison tasks: Quantitative and qualitative measurements. *Advances in Computational Intelligence*, 4, Article 7.
11. Gergelyfi, M., Jacob, B., Olivier, E., & Zénon, A. (2015). Dissociation between mental fatigue and motivational state during prolonged mental effort. *Frontiers in Human Neuroscience*, 9, 176.
12. Grandjean, E. (2000). *Fitting the task to the man: A textbook of occupational ergonomics*. Taylor & Francis.
13. Grobelny, J., Glinka, M., & Chirkowska-Smolak, T. (2024). The impact of hedonic social media use during microbreaks on employee resources recovery. *Scientific Reports*, 14, 21603.
14. Hockey, G. R. J. (2013). *The psychology of fatigue: Work, effort and control*. Cambridge University Press.
15. Hofmann, W., Schmeichel, B. J., & Baddeley, A. D. (2012). Executive functions and self-regulation. *Trends in Cognitive Sciences*, 16(3), 174–180. <https://doi.org/10.1016/j.tics.2012.01.006>
16. Hunter, E. M., & Wu, C. (2016). Give me a better break: Choosing workday break activities to maximize resource recovery. *Journal of Applied Psychology*, 101(2), 302–311.
17. Inzlicht, M., Schmeichel, B. J., & Macrae, C. N. (2014). Why self-control seems limited: A resource depletion perspective. *Trends in Cognitive Sciences*, 18(3), 127–133.
18. Kaplan, S. (2001). Meditation, restoration, and the management of mental fatigue. *Environment and Behavior*, 33(4), 480–506.
19. Kellogg, S., & Kun, A. L. (2019). Cognitive workload and human performance in sustained attention tasks. *Human Factors*, 61(7), 1121–1135.
20. Lim, J., & Dinges, D. F. (2010). A meta-analysis of the impact of short-term sleep deprivation on cognitive variables. *Psychological Bulletin*, 136(3), 375–389.

21. Lorist, M. M., Boksem, M. A. S., & Ridderinkhof, K. R. (2005). Impaired cognitive control and reduced efficiency during mental fatigue. *Cognitive Brain Research*, 24(2), 199–205.
22. Lyubych, Z., Gulseren, D., Premji, Z., Wingate, T. G., Deng, C., Bélanger, L. J., & Turner, N. (2022). Role of work breaks in well-being and performance: A systematic review and future research agenda. *Journal of Occupational Health Psychology*, 27(5), 470–487.
23. Marcora, S. M., Staiano, W., & Manning, V. (2009). Mental fatigue impairs physical performance. *Journal of Applied Physiology*, 106(3), 857–864.
24. Meijman, T. F., & Mulder, G. (2000). Psychological aspects of workload. In P. J. D. Drenth & H. Thierry (Eds.), *Handbook of work and organizational psychology* (pp. 5–33). Psychology Press.
25. Muraven, M., & Baumeister, R. F. (2000). Self-regulation and depletion of limited resources. *Psychological Bulletin*, 126(2), 247–259.
26. Pashler, H., Johnston, J. C., & Ruthruff, E. (2001). Attention and performance. *Annual Review of Psychology*, 52, 629–651.
27. Persson, J., & Reuter-Lorenz, P. A. (2008). Gaining control: Training executive function and working memory. *Psychological Science*, 19(11), 1123–1129.
28. Randolph, J. J. (2012). The psychology of sustained attention and cognitive recovery. *Educational Psychology Review*, 24, 445–463.
29. Saito, H., Lee, A., Bui, H. T., & Baum, T. (2024). Fostering employee well-being and workplace recovery in the hospitality industry: The role of supportive break times at work. *Journal of Hospitality and Tourism Management*, 61, 91–100.
30. Scholey, A. B., Harper, D. J., & Kennedy, D. O. (2001). Cognitive demand and fatigue effects on performance. *Psychopharmacology*, 154, 271–278.
31. Sharma, P., Justus, J. C., Thapa, M., & Poudel, G. R. (2023). Sensors and systems for monitoring mental fatigue: A systematic review. *IEEE/related technical literature review sources*.
32. Smallwood, J., & Schooler, J. W. (2015). The science of mind wandering. *Annual Review of Psychology*, 66, 487–518.
33. Sonnentag, S., & Fritz, C. (2015). Recovery from job stress: The stressor-detachment model. *Journal of Organizational Behavior*, 36(S1), S72–S103.

34. Taris, T. W., & Kompier, M. A. J. (2005). Job demands, job control, and fatigue. *Work & Stress*, 19(2), 1–12.
35. Trougakos, J. P., Beal, D. J., Green, S. G., & Weiss, H. M. (2008). Making the break count: An episodic examination of recovery activities, emotional experiences, and positive affect during work breaks. *Journal of Applied Psychology*, 93(6), 1317–1331.
36. Westbrook, A., & Braver, T. S. (2016). Cognitive effort: A neuroeconomic approach. *Cognitive, Affective, & Behavioral Neuroscience*, 16, 395–415.
37. Zacher, H., & Rudolph, C. W. (2021). Individual differences and micro-break recovery experiences at work. *Occupational Health Psychology Research*.